

General monotonicity, interpolation of operators, and applications.

S.M. Grigoriev, Y. Sagher, T.R. Savage

1. Introduction.

Definition 1.1. A measurable function f on (Ω, Σ, μ) is called *admissible*, if the following conditions hold:

- (1) $\{f \neq 0\}$ is a σ -finite set.
- (2) $f < \infty$ a.e. on Ω .
- (3) $\mu(\{x : |f(x)| > \gamma\}) < \infty$ for some $\gamma > 0$.

Recall the definition of the decreasing rearrangement of a function.

Definition 1.2. Let (Ω, Σ, μ) be a measure space and assume that f is an admissible function on Ω . Define, for each $\alpha \geq 0$, the *distribution function* of f : $f_*(\alpha) = \mu(\{x : |f(x)| > \alpha\})$.

Definition 1.3. For each $x \geq 0$, define

$$f^*(x) = \begin{cases} \inf(\{\alpha : f_*(\alpha) \leq x\}) & \text{if } \exists \alpha : f_*(\alpha) \leq x \\ \infty & \text{otherwise} \end{cases}. \quad (1)$$

f^* is called the *decreasing rearrangement* of f .

The distribution function of f^* with respect to Lebesgue measure on $\mathbb{R}^+$ is equal to the distribution function of f on (Ω, Σ, μ) :

Theorem 1.4. $(f^*)_*(\alpha) = f_*(\alpha), \forall \alpha \geq 0$.

Definition 1.5. Considering sequences as functions defined on $\Omega \subseteq \mathbb{Z}$, with $\mu = \#$, the counting measure, we use formula (1) to define for each $k \in \mathbb{N}$,

$$a_k^* = (\{a_n\}^*)(k-1). \quad (2)$$

We recall the definition of weighted L^q -norms.

Definition 1.6. Let $0 < q \leq \infty$, and let w be a nonnegative measurable function on (Ω, Σ, μ) . Assume that f is a measurable function on (Ω, Σ, μ) . Define the *weighted L^q -norm* of f with the weight w :

$$\|f\|_{L_w^q(\Omega, \Sigma, \mu)} = \|f \cdot w\|_{L^q(\Omega, \Sigma, \mu)}.$$

$L_w^q = L_w^q(\Omega, \Sigma, \mu)$ is the space of all measurable functions f for which $\|f\|_{L_w^q(\Omega, \Sigma, \mu)}$ is finite.

In particular, working with $(\mathbb{N}, 2^{\mathbb{N}}, \#)$, we follow the practice of writing

$$\|\{a_n\}\|_{l_w^q} = \|\{a_n \cdot w(n)\}\|_{l^q}.$$

l_w^q is the space of sequences $\{a_n\}$ for which $\|\{a_n\}\|_{l_w^q}$ is finite.

We will make extensive use of the spaces of functions defined on $\mathbb{R}^+$ with the following weight:

Definition 1.7. Let $0 < p \leq \infty$, $0 < q \leq \infty$. We denote, $\forall x > 0$:

$$w(p, q)(x) = x^{\frac{1}{p} - \frac{1}{q}}. \quad (3)$$

When we work with sequences, we denote, $\forall n \geq 1$:

$$w(p, q)(n) = n^{\frac{1}{p} - \frac{1}{q}}. \quad (4)$$

Definition 1.8.

(i) Assume that f is a measurable function on (Ω, Σ, μ) and that $f^* < \infty$ on $\mathbb{R}^+$. For $0 < p < \infty$, $0 < q \leq \infty$, and $p = q = \infty$, define the $L(p, q)$ -norm of f :

$$\|f\|_{L(p, q)} = \|f\|_{L(p, q)(\Omega, \Sigma, \mu)} = \|f^*\|_{L_{w(p, q)}^q(\mathbb{R}^+, \mathcal{B}, \lambda)} \quad (5)$$

$L(p, q) = L(p, q)(\Omega, \Sigma, \mu)$ is the space of all measurable functions f , for which $\|f\|_{L(p, q)}$ is finite.

(ii) Analogously,

$$\|\{a_n\}\|_{l(p, q)} = \|\{a_n^*\}\|_{l_{w(p, q)}^q}. \quad (6)$$

$l(p, q)$ is the collection of all sequences $\{a_n\}$ for which $\|\{a_n\}\|_{l(p, q)}$ is finite.

Lemma 1.9. Assume that $\{a_n\}^*$ exists. For $0 < p, q \leq \infty$,

$$\|\{a_k\}\|_{l(p, q)} \sim \|\{a_n\}^*(\cdot)\|_{L_{w(p, q)}^q}. \quad (7)$$

Definition 1.10. ONB is the collection of is an orthonormal uniformly bounded sequences $\{\phi_n\}$ of complex-valued measurable functions.

Theorem 1.11 (R.E.A.C. Paley, [10]).

Let $\{\phi_n\}$ be an ONB on (Ω, Σ, μ) , $1 < p \leq 2$, and $c_n = \int_{\Omega} f \bar{\phi}_n d\mu$:

(i) If $f \in L^p(\Omega, \Sigma, \mu)$, then $\|\{c_n\}\|_{l(p', p)} \leq B \|f\|_{L^p}$.

(ii) If $\|\{c_n\}\|_{l(p, p')} < \infty$ then $\|f\|_{L^p} \leq B' \|\{c_n\}\|_{l(p, p')}$.

The following classical theorem shows that, in the context of $L(p, q)$ spaces, Paley's theorem is best possible.

Theorem 1.12 (G.H. Hardy and J.E. Littlewood, [6]).

Let $\{c_n\} \searrow 0$, $f(x) = \sum_{n=0}^{\infty} c_n \cos nx$ or $f(x) = \sum_{n=1}^{\infty} c_n \sin nx$, $p > 1$. Then

$$\|f\|_{L^p(0, \pi)} \sim \|\{c_n\}\|_{l(p', p)}.$$

Hardy and Littlewood also described a norm relation between monotone decreasing functions and their trigonometric Fourier coefficients:

Theorem 1.13 (G.H. Hardy and J.E. Littlewood, [7]).

Let $f \searrow$ on $[0, \pi]$, $f \geq 0$, $1 < p < \infty$, and let $\{c_n\}$ be the sequence of trigonometric Fourier coefficients of f . Then:

$$\|f\|_{L^p(0, \pi)} \sim |c_0| + \|\{c_n\}\|_{l_{w(p', p)}^p}.$$

The extension of Theorem 1.13 to broader classes of sequences started with the introduction of *quasi-monotone sequences* by Shah [13]; this concept also applies to functions defined on $(0, \infty)$:

Definition 1.14. Let $\beta > 0$.

(a) If $a_n \geq 0$ and $\{a_n \cdot n^{-\beta}\} \searrow$, the sequence $\{a_n\}$ is called a β -*quasi-monotone decreasing sequence* (written as $\{a_n\} \in QDS(\beta)$).

(b) If $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is such that $x^{-\beta}f(x) \searrow$ then f is called a β -*quasi-monotone decreasing function*, $f \in QD(\beta)$.

Denote $QDS = \bigcup_{\beta>0} QDS(\beta)$; $QD = \bigcup_{\beta>0} QD(\beta)$. We say that $\{a_n\}$ is a *quasi-monotone decreasing sequence* if $\{a_n\} \in QDS$ and that f is a *quasi-monotone decreasing function* if $f \in QD$.

The interest in quasi-monotone sequences stems from the fact that the class of trigonometric series with quasi-monotone Fourier coefficients is closed under term-by-term differentiation. For an application, see [11].

Askey and Wainger [1] proved that if $\{c_n\}$ is a quasi-monotone sequence of trigonometric Fourier coefficients of f then for $1 < p < \infty$, $1 \leq q < \infty$,

$$\|f\|_{L^q_{w(p,q)(0,\pi)}} \sim \|\{c_n\}\|_{l^q_{w(p',q)}}. \quad (8)$$

An extension of Real Interpolation theory made it possible to simplify the proof of (8). The extension consists of interpolation of normed *monoids* rather than normed vector spaces.

Tikhonov [14] defined the following extension of the class of quasi-monotone sequences:

Definition 1.15. Let $B \geq 1$. We write $\{a_n\} \in GMS(B)$ if for all $n \geq 1$:

$$\sum_{k=n}^{2n-1} |a_k - a_{k+1}| \leq B |a_n|. \quad (9)$$

Define $GMS = \bigcup_{B \geq 1} GMS(B)$. If $\{a_n\} \in GMS$, we say that $\{a_n\}$ is a *general monotone sequence*.

Liflyand and Tikhonov [9] defined *general monotone functions* (GM). We will work with a special case of their definition.

Definition 1.16. Let $B \geq 1$. We write $f \in GM(B)$ if $f \in BV_{loc}((0, \infty))$ and for all $x > 0$:

$$V_f([x, 2x]) \leq B |f(x)|, \quad (10)$$

where $V_f([a, b])$ is the variation of f on $[a, b]$. Define $GM = \bigcup_{B \geq 1} GM(B)$.

If $f \in GM$, we say that f is a *general monotone function*.

We will often use the following classes of functions and sequences:

Definition 1.17.

$$GM_1(B) = \{f \in \mathcal{M}_{\mathbb{R}^+} : |f(t)| \leq B |f(x)|, \forall x \leq t \leq 2x\}; \quad (11)$$

$$GMS_1(B) = \{\{a_k\}_{k=1}^{\infty} : |a_k| \leq B |a_n|, \forall n \leq k \leq 2n\}; \quad (12)$$

$$GM_2(B) = \left\{ f \in BV_{loc}(\mathbb{R}^+) : V_f([x, M]) \leq B \left(|f(x)| + \int_x^M |f(t)| \frac{dt}{t} \right), \forall M > 0, \forall 0 < x < M \right\}, \quad (13)$$

$$GMS_2(B) = \left\{ \{a_k\}_{k=1}^\infty : \sum_{k=n}^{N-1} |a_k - a_{k+1}| \leq B \left(|a_n| + \sum_{k=n+1}^N \frac{|a_k|}{k} \right), \forall N > 1, \forall 1 \leq n < N \right\}. \quad (14)$$

We write for $j = 1, 2$, $GM_j = \bigcup_{B \geq 1} GM_j(B)$, $GM_j^+ = \{f \in GM_j : f \geq 0\}$, $GMS_j = \bigcup_{B \geq 1} GMS_j(B)$, and $GMS_j^+ = GMS_j \cap \{\{a_n\} : a_n \geq 0, \forall n\}$.

Definition 1.18. For each $0 \leq \varphi < \frac{\pi}{2}$ and $\alpha \in \mathbb{R}$, define

$$S_{\alpha, \varphi} := \{z \in \mathbb{C} : |\arg(e^{-i\alpha} z)| \leq \varphi\} \cup \{0\}. \quad (15)$$

Lemma 1.19. Let (Ω, Σ, μ) be a measure space and assume that $f(\omega) \in S_{\alpha, \varphi}$ for a.e. $\omega \in \Omega$. Then

$$\int_{\Omega} |f| d\mu \leq \frac{1}{\cos \varphi} \left| \int_{\Omega} f d\mu \right|. \quad (16)$$

Definition 1.20. For each $0 \leq \varphi < \frac{\pi}{2}$ and $0 \leq \alpha < 2\pi$, define

$$GM_{\alpha, \varphi} = \{f \in GM : f(x) \in S_{\alpha, \varphi}, \mu\text{-a.e. } x > 0\};$$

$$GMS_{\alpha, \varphi} = \{\{a_n\} \in GMS : a_n \in S_{\alpha, \varphi}, \forall n \geq 1\}.$$

We write $GM_{\alpha, \varphi}(B) = GM_{\alpha, \varphi} \cap GM(B)$ and $GMS_{\alpha, \varphi}(B) = GMS_{\alpha, \varphi} \cap GMS(B)$.

Remark 1.21. Observe that $GMS_{0,0} = \{\{a_n\} \in GMS : a_n \geq 0\} =: GMS^+$, $GM_{0,0} = \{f \in GM : f \geq 0\} =: GM^+$.

Lemma 1.22 (E. Liflyand and S. Tikhonov, [9]).

$$GM(B) \subset GM_1(2B) \cap GM_2(2B^2); \quad (17)$$

$$GM_1(B) \cap GM_2(B) \subset GM(2B^2). \quad (18)$$

Analogously for general monotone sequences:

Lemma 1.23 (S. Tikhonov, [14]).

$$GMS(B) \subset GMS_1(2B) \cap GMS_2(2B^2); \quad (19)$$

$$GMS_1(B) \cap GMS_2(B) \subset GMS(2B^2). \quad (20)$$

2. Some results from Real Interpolation theory.

Definition 2.1. A *semigroup* in a topological vector space (t.v.s.) is a subset G of the t.v.s. that is closed under addition. A semigroup that contains 0 is a *monoid*. A monoid that is closed under multiplication by any positive scalar is a *cone*.

Definition 2.2. A *quasi-norm* on a vector space X is a function $\|\cdot\|_X : X \rightarrow [0, \infty)$ that satisfies:

- (a) $\|a\|_X = 0 \Leftrightarrow a = 0$.
- (b) $\exists k = k(X)$ such that $\forall a_1, a_2 \in X: \|a_1 + a_2\|_X \leq k(\|a_1\|_X + \|a_2\|_X)$.
- (c) $\|\lambda a\|_X = |\lambda| \|a\|_X, \forall \lambda \in \mathbb{C}, a \in X$.

A vector space equipped with a quasi-norm is a *quasi-normed vector space*, a complete quasi-normed vector space is called a *quasi-Banach space*.

Definition 2.3. If G_0 and G_1 are monoids continuously embedded in a t.v.s., we say that (G_0, G_1) is an *interpolation couple*.

Throughout this section, G_0, G_1 will be quasi-normed monoids which form an interpolation couple in a t.v.s. of measurable functions, the topology of the t.v.s. is defined by the convergence in measure.

Definition 2.4. For $f \in G_0 + G_1, t > 0$, define

$$K(t, f) = K(t, f, G_0, G_1) \\ = \inf \{ \|f_0\|_{G_0} + t \|f_1\|_{G_1} \mid f_0 + f_1 = f, f_j \in G_j, j = 0, 1 \}. \quad (21)$$

Definition 2.5. (K -method of interpolation of monoids).

For $f \in G_0 + G_1$, and for each $0 < \theta < 1$ and $0 < q \leq \infty$ or for $0 \leq \theta \leq 1$ and $q = \infty$, define

$$\|f\|_{(G_0, G_1)_{\theta, q; K}} = \|t^{-\theta} K(t, f)\|_{L^q((0, \infty), \frac{dt}{t})}. \quad (22)$$

Define the *Interpolation monoid with respect to K* as

$$(G_0, G_1)_{\theta, q; K} = \left\{ f \in G_0 + G_1 \mid \|f\|_{(G_0, G_1)_{\theta, q; K}} < \infty \right\}. \quad (23)$$

It is clear that $(G_0, G_1)_{\theta, q; K}$ is a monoid. When G_0 and G_1 are quasi-normed vector spaces, we call $(G_0, G_1)_{\theta, q; K}$ an *Interpolation space*.

Theorem 2.6 (J. Gilbert, [5]).

Assume X is a quasi-normed monoid, w is a measurable positive function on Ω , σ is positive, piecewise continuous function such that both σ and σ_t , where $\sigma_t(\lambda) := t\lambda\sigma(t\lambda)$, are in $L^\infty(0, \infty), \forall t > 0$. For any $f \in (X, X_w)_{\theta, q; K}, \theta \in (0, 1)$, and $q \in (0, \infty]$,

$$\|f\|_{\theta, q; K} \sim \left(\int_0^\infty (t^{-\theta} \|f \cdot \sigma_t \circ w\|_X)^q \frac{dt}{t} \right)^{1/q}. \quad (24)$$

3. Selected results from the theory of $L(p, q)$ spaces.

Theorem 3.1 ([8]). Suppose that I is a subinterval of $\mathbb{Z}$ and that $\{a_k\}_{k \in I}$, $\{c_k\}_{k \in I}$ are nonnegative sequences. If $\{c_k\} \searrow$ then $\sum_{k \in I} a_k c_k \leq \sum_{k \in I} a_k^* c_k$. If $\{c_k\} \nearrow$ then $\sum_{k \in I} a_k c_k \geq \sum_{k \in I} a_k^* c_k$.

Theorem 3.2 (G.H. Hardy).

Let f_1, f_2, g be nonnegative, measurable functions on $\mathbb{R}^+$.

(a) Assume that $g \searrow$ and that for all $a > 0$, $\int_0^a f_1 dt \leq \int_0^a f_2 dt$. Then

$$\int_0^\infty f_1 g dt \leq \int_0^\infty f_2 g dt. \quad (25)$$

(b) Assume that $g \nearrow$ and that for all $a > 0$, $\int_a^\infty f_1 dt \leq \int_a^\infty f_2 dt$. Then (25) holds.

Corollary 3.3. Suppose that f, g are nonnegative, measurable on $\mathbb{R}^+$ functions.

If $g \searrow$ then $\int_0^\infty f g dt \leq \int_0^\infty f^* g dt$. If $g \nearrow$ then $\int_0^\infty f g dt \geq \int_0^\infty f^* g dt$.

Theorem 3.4. Let f be measurable on $\mathbb{R}^+$. For $0 < p \leq q \leq \infty$,

$$\|f\|_{L(p,q)(0,\infty)} \leq \|f\|_{L_{w(p,q)}^q(0,\infty)}. \quad (26)$$

The following theorem is proved for sine Fourier coefficients in [12].

Theorem 3.5. Let $f \in L(p', q)$, $1 < p < \infty$, $0 < q \leq \infty$ and assume that $\{c_n\}$ are Fourier coefficients of f with respect to $\{e^{int}\}_{n=-\infty}^\infty$. Define

$\sigma_n = \frac{1}{2n+1} \sum_{k=-n}^n c_k$, and let $\{m(a_n)\}$ be the least bell-shaped majorant of $\{a_n\}$, that is to say, $m(a_n) := \sup \{|a_k| : |k| \geq |n|\}$. Then $\exists A(p) > 0$ so that:

$$\|\{m(\sigma_n)\}\|_{l(p,q)} \leq A(p) \|f\|_{L(p',q)(0,2\pi)}, \quad (27)$$

Consequently,

$$\|\{\sigma_n\}\|_{l(p,q)} \leq A(p) \|f\|_{L(p',q)(0,2\pi)}. \quad (28)$$

4. Some elementary properties of GM and GMS .

Lemma 4.1. For $j = 1, 2$, if $f \in GM_j$ then $\{f(n)\} \in GMS_j$. Moreover, if $\{a_n\} \in GMS_1$ and $f(x) = a_{\lceil x \rceil}$ then $f \in GM_1$.

Lemma 4.2. If $f \in GM$ then $\{f(n)\} \in GMS$. Conversely, for each $\{a_n\} \in GMS$ there is an $f \in GM$ such that $f(n) = a_n$, $\forall n \geq 1$.

Using Lemma 4.2, results for GMS follow from corresponding results for GM , and therefore we shall only give proofs for GM functions.

Lemma 4.3. Assume that $f \in GM_1(B)$ and that f^* exists. Then $\forall x > 0$,

$$|f(x)| \leq Bf^*\left(\left(\frac{x}{2}\right) -\right). \quad (29)$$

Proof.

Let $x > 0$, $0 < \theta < 1$. Since (29) trivially holds if $f(x) = 0$, we can assume that $f(x) \neq 0$. For all $t \in [\frac{x}{2}, x]$, $t \leq x \leq 2t$, and by (11),

$$|f(t)| \geq \frac{|f(x)|}{B} > \frac{\theta |f(x)|}{B}.$$

Therefore:

$$f_*\left(\frac{\theta |f(x)|}{B}\right) = \lambda \left\{ t : |f(t)| > \frac{\theta |f(x)|}{B} \right\} \geq \frac{x}{2} > \frac{\theta x}{2},$$

and since $f_* \searrow$:

$$f^*\left(\frac{\theta x}{2}\right) = \inf \left\{ \alpha : f_*(\alpha) \leq \frac{\theta x}{2} \right\} \geq \frac{\theta |f(x)|}{B},$$

or $|f(x)| \leq \frac{B}{\theta} f^*\left(\frac{\theta x}{2}\right)$. Letting $\theta \rightarrow 1^-$, (29) follows. $\square$

The corresponding result for GMS_1 was proved in [3]:

Lemma 4.4 (B. Booton, [3]).

Let $\{c_k\} \in GMS_1(B)$ be such that $\{c_k^*\}$ exists. Then $\forall n \geq 1$,

$$|c_n| \leq Bc_{\lfloor \frac{n}{2} \rfloor + 1}^*.$$

Lemma 4.5. Assume that $f \in GM_1$, $A > 0$, $\alpha \in \mathbb{R}$.

(i) If $\int_A^\infty |f(x)| x^\alpha dx < \infty$ then $\lim_{x \rightarrow \infty} x^{\alpha+1} f(x) = 0$.

(ii) If $\int_0^A |f(x)| x^\alpha dx < \infty$ then $\lim_{x \rightarrow 0+} x^{\alpha+1} f(x) = 0$.

Proof.

Let $2^k \leq x \leq 2^{k+1}$. If $\alpha < 0$ then $2^\alpha \cdot 2^{k\alpha} = 2^{(k+1)\alpha} \leq x^\alpha \leq 2^{k\alpha}$. If $\alpha \geq 0$ then $2^{k\alpha} \leq x^\alpha \leq 2^{(k+1)\alpha} = 2^\alpha \cdot 2^{k\alpha}$. That is to say, for any $\alpha \in \mathbb{R}$:

$$\min\{2^\alpha, 1\} \cdot 2^{k\alpha} \leq x^\alpha \leq \max\{2^\alpha, 1\} \cdot 2^{k\alpha}.$$

Furthermore, by (11), $\frac{1}{B} |f(2^{k+1})| \leq |f(x)| \leq B |f(2^k)|$. Therefore:

$$\frac{1}{B} \min \{2^\alpha, 1\} 2^{k\alpha} |f(2^{k+1})| \leq x^\alpha |f(x)| \leq B \max \{2^\alpha, 1\} 2^{k\alpha} |f(2^k)|. \quad (30)$$

It follows from (30) that for all $-\infty \leq n < N \leq \infty$,

$$\begin{aligned} \int_{2^n}^{2^N} |f(x)| x^\alpha dx &= \sum_{k=n}^{N-1} \int_{2^k}^{2^{k+1}} x^\alpha |f(x)| dx \\ &\geq \frac{\min \{2^\alpha, 1\}}{B} \cdot \sum_{k=n}^{N-1} 2^{k\alpha} |f(2^{k+1})| \cdot 2^k = \frac{\min \{2^\alpha, 1\}}{B \cdot 2^{\alpha+1}} \sum_{k=n+1}^N 2^{k(\alpha+1)} |f(2^k)|. \end{aligned}$$

Therefore, if $\int_A^\infty |f(x)| x^\alpha dx < \infty$ then $\sum_{k=\lceil \log_2 A \rceil}^\infty 2^{k(\alpha+1)} |f(2^k)| < \infty$, and so

$\lim_{k \rightarrow \infty} 2^{k(\alpha+1)} |f(2^k)| = 0$. If $\int_0^A |f(x)| x^\alpha dx < \infty$ then $\sum_{k=-\infty}^{\lfloor \log_2 A \rfloor} 2^{k(\alpha+1)} |f(2^k)| < \infty$, and so $\lim_{k \rightarrow -\infty} 2^{k(\alpha+1)} |f(2^k)| = 0$. The claim of the Lemma now follows by substituting $\alpha + 1$ for α in the right-hand side of (30). $\square$

Corollary 4.5. If $0 < p \leq \infty$, $0 < q < \infty$ and $f \in L_{w(p,q)}^q \cap GM_1$ then $\lim_{x \rightarrow 0+} x^{\frac{1}{p}} f(x) = \lim_{x \rightarrow \infty} x^{\frac{1}{p}} f(x) = 0$.
By Lemma 4.1:

Lemma 4.6. Assume $\{a_n\} \in GMS_1$. If $\alpha \in \mathbb{R}$ is such that $\sum_{n=1}^\infty n^\alpha |a_n| < \infty$ then $\lim_{n \rightarrow \infty} n^{\alpha+1} a_n = 0$.

Corollary 4.6. If $0 < p \leq \infty$, $0 < q < \infty$ and $\{a_n\} \in l_{w(p,q)}^q \cap GMS_1$ then $\lim_{n \rightarrow \infty} n^{\frac{1}{p}} a_n = 0$.

If $\alpha = -1$ then the claim of Lemma 4.6 is also true for GMS_2 :

Lemma 4.7. Assume $\{a_n\} \in l_{\frac{1}{k}}^1 \cap GMS_2$. Then $\lim_{n \rightarrow \infty} a_n = 0$.

Proof.

By the assumption of the Lemma, $\sum_{m=1}^\infty |a_m - a_{m+1}| \leq B \sum_{k=1}^\infty \frac{|a_k|}{k} < \infty$, and so, $\lim_{m \rightarrow \infty} (a_1 - a_m) = \sum_{j=1}^\infty (a_m - a_{m+1})$ exists and finite. If $0 \neq a = \lim_{m \rightarrow \infty} a_m$ then $\exists N$ such that $||a_m| - |a|| < \frac{|a|}{2}$, $\forall m \geq N$, that is, $\frac{|a|}{2m} \leq \frac{|a_m|}{m}$. But then $\sum_{m=N}^\infty \frac{|a|}{2m} < \infty$, since $\{a_n\} \in l_{\frac{1}{k}}^1$, a contradiction. $\square$

Lemma 4.8. Let $B \geq 1$, $f \in GM_1(B)$. Then for $0 < p < \infty$, $0 < q < \infty$:

$$\frac{C_1(p, q)}{B} \|f\|_{L_{w(p, q)}^q} \leq \left(\sum_{k=-\infty}^{\infty} 2^{\frac{kq}{p}} |f(2^k)|^q \right)^{\frac{1}{q}} \leq BC_2(p, q) \|f\|_{L(p, q)}, \quad (31)$$

with $0 < C_j(p, q) < \infty$. For $0 < q < \infty$,

$$\|f\|_{L_{w(\infty, q)}^q} = \left(\int_0^{\infty} |f(x)|^q \frac{dx}{x} \right)^{\frac{1}{q}} \leq B (\ln 2)^{\frac{1}{q}} \left(\sum_{k=-\infty}^{\infty} |f(2^k)|^q \right)^{\frac{1}{q}}. \quad (32)$$

Furthermore, for $0 < p \leq \infty$:

$$\begin{aligned} \frac{1}{2^{\frac{1}{p}} B} \|f\|_{L(p, \infty)} &\leq \frac{1}{2^{\frac{1}{p}} B} \|f\|_{L_{w(p, \infty)}^{\infty}} \leq \sup_{k \in \mathbb{Z}} \left\{ 2^{\frac{k}{p}} |f(2^k)| \right\} \\ &\leq 2^{\frac{1}{p}} B \|f\|_{L_{w(p, \infty)}^{\infty}} \leq 2^{\frac{2}{p}} B^2 \|f\|_{L(p, \infty)}. \end{aligned} \quad (33)$$

Proof.

For $f \in GM_1(B)$ and $2^k \leq x \leq 2^{k+1}$, $|f(x)| \leq B |f(2^k)|$, and so:

$$\begin{aligned} \|f\|_{L_{w(p, q)}^q}^q &= \int_0^{\infty} x^{\frac{q}{p}-1} |f(x)|^q dx = 2^{\frac{q}{p}} \sum_{k=-\infty}^{\infty} \int_{2^k}^{2^{k+1}} \left(\frac{x}{2} \right)^{\frac{q}{p}} |f(x)|^q \frac{dx}{x} \\ &\leq 2^{\frac{q}{p}} B^q \sum_{k=-\infty}^{\infty} 2^{\frac{kq}{p}} |f(2^k)|^q \int_{2^k}^{2^{k+1}} \frac{dx}{x} = 2^{\frac{q}{p}} B^q \ln 2 \sum_{k=-\infty}^{\infty} 2^{\frac{kq}{p}} |f(2^k)|^q, \end{aligned}$$

proving the first inequality in (31) and (32). Assume $2^{k-2} \leq x \leq 2^{k-1}$, or equivalently, $2x \leq 2^k \leq 4x$. If $q > p$ then $2^{k(\frac{q}{p}-1)} \leq (4x)^{\frac{q}{p}-1}$, if $q \leq p$ then $2^{k(\frac{q}{p}-1)} \leq (2x)^{\frac{q}{p}-1}$ and so, for all $0 < p, q < \infty$:

$$2^{k(\frac{q}{p}-1)} \leq A(p, q) x^{\frac{q}{p}-1},$$

$A(p, q) = \max \left\{ 2^{\frac{q}{p}-1}, 4^{\frac{q}{p}-1} \right\}$. By Lemma 4.3, $|f(2^k)| \leq B f^*(2^{k-1}-)$, and so:

$$\begin{aligned} \sum_{k=-\infty}^{\infty} 2^{\frac{kq}{p}} |f(2^k)|^q &\leq B^q \sum_{k=-\infty}^{\infty} 2^{\frac{kq}{p}} (f^*(2^{k-1}-))^q \\ &= 4B^q \sum_{k=-\infty}^{\infty} \int_{2^{k-2}}^{2^{k-1}} 2^{k(\frac{q}{p}-1)} (f^*(2^{k-1}-))^q dx \leq 4A(p, q) B^q \int_0^{\infty} x^{\frac{q}{p}-1} (f^*(x))^q dx \end{aligned}$$

$$= B^q A(p, q) \int_0^\infty x^{\frac{q}{p}-1} (f^*(x))^q dx = B^q A(p, q) \|f\|_{L(p, q)}^q,$$

proving the second inequality in (31).

Let us consider (33). Observe that $|f(x)| \leq x^{-\frac{1}{p}} \sup_{x>0} \left\{ x^{\frac{1}{p}} |f(x)| \right\}$, and so, $|f^*(x)| \leq x^{-\frac{1}{p}} \sup_{x>0} \left\{ x^{\frac{1}{p}} |f(x)| \right\}$, that is to say, $\|f\|_{L(p, \infty)} \leq \|f\|_{L_w^\infty(p, \infty)}$, proving the first inequality. For $2^k \leq x \leq 2^{k+1}$ and $\alpha = \frac{1}{p}$, (30) shows that:

$$\frac{1}{B} \cdot 2^{\frac{k}{p}} |f(2^{k+1})| \leq x^{\frac{1}{p}} |f(x)| \leq 2^{\frac{1}{p}} B \cdot 2^{\frac{k}{p}} |f(2^k)|,$$

or equivalently, using $2^{k-1} \leq \frac{x}{2} \leq 2^k$:

$$\frac{1}{2^{\frac{1}{p}} B} x^{\frac{1}{p}} |f(x)| \leq 2^{\frac{k}{p}} |f(2^k)| \leq 2^{\frac{1}{p}} B \cdot \left(\frac{x}{2}\right)^{\frac{1}{p}} \left|f\left(\frac{x}{2}\right)\right|.$$

Taking the supremum over all $k \in \mathbb{Z}$ we obtain:

$$\frac{1}{2^{\frac{1}{p}} B} \|f\|_{L_w^\infty(p, \infty)} \leq \sup_{k \in \mathbb{Z}} \left\{ 2^{\frac{k}{p}} |f(2^k)| \right\} \leq 2^{\frac{1}{p}} B \|f\|_{L(p, \infty)},$$

proving the second and the third inequality. Finally, by Lemma 4.3, for all $x > 0$, $|f(x)| \leq B f^*\left(\left(\frac{x}{2}\right) -\right)$, therefore, for all $0 < \theta < 1$:

$$\begin{aligned} \sup_{x>0} \left\{ x^{\frac{1}{p}} |f(x)| \right\} &\leq B \sup_{x>0} \left\{ x^{\frac{1}{p}} f^*\left(\left(\frac{x}{2}\right) -\right) \right\} = 2^{\frac{1}{p}} B \sup_{x>0} \left\{ x^{\frac{1}{p}} f^*(x-) \right\} \\ &\leq 2^{\frac{1}{p}} B \sup_{x>0} \left\{ x^{\frac{1}{p}} f^*(\theta x) \right\} = \frac{2^{\frac{1}{p}} B}{\theta^{\frac{1}{p}}} \sup_{x>0} \left\{ (\theta x)^{\frac{1}{p}} f^*(\theta x) \right\} = \frac{2^{\frac{1}{p}} B}{\theta^{\frac{1}{p}}} \sup_{x>0} \left\{ x^{\frac{1}{p}} f^*(x) \right\}. \end{aligned}$$

Letting $\theta \rightarrow 1-$, it follows that $\|f\|_{L_w^\infty(p, \infty)} \leq 2^{\frac{1}{p}} B \|f\|_{L(p, \infty)}$, and the last inequality in (33) follows. $\square$

The next lemma shows that the statement of Theorem 3.4 holds for $q < p$ if $f \in GM_1$. The proof is similar to the proof by B. Booton [3] of the analogous statement for GMS_1 .

Lemma 4.9. Let $f \in GM_1(B)$, $0 < q < p$. Then

$$\|f\|_{L(p, q)} \leq \left(\frac{2p}{q}\right)^{\frac{1}{q}} B^2 \|f\|_{L_w^q(p, q)}.$$

Proof.

Let $g(x) = \sum_{k=-\infty}^{\infty} |f(2^k)| I_{[2^k, 2^{k+1})}(x)$. Then by (11), $|f(x)| \leq Bg(x)$. Furthermore, g^* is decreasing, right-continuous and for each $x > 0$, $g^*(x) = |f(2^k)|$ for some $k \in \mathbb{Z}$. Therefore, g^* can be written as

$$g^*(x) = \sum_{k \in J \cap \mathbb{Z}} \alpha_k I_{[x_k, x_{k+1})}(x),$$

where J is an interval, $\alpha_k > \alpha_{k+1}$, and $[x_k, x_{k+1}) = \{g^* = \alpha_k\}$, $\forall k$. And so:

$$\begin{aligned} \|f\|_{L(p,q)}^q &\leq B^q \|g\|_{L(p,q)}^q = B^q \int_0^\infty x^{\frac{q}{p}} (g^*(x))^q \frac{dx}{x} \\ &= B^q \sum_{k=-\infty}^{\infty} \int_{\{g^*=\alpha_k\}} x^{\frac{q}{p}} \alpha_k^q \frac{dx}{x} = B^q \sum_{k=-\infty}^{\infty} \alpha_k^q \int_{x_k}^{x_{k+1}} x^{\frac{q}{p}} \frac{dx}{x} \\ &= B^q \frac{p}{q} \sum_{k=-\infty}^{\infty} \alpha_k^q \left(x_{k+1}^{\frac{q}{p}} - x_k^{\frac{q}{p}} \right) \leq B^q \frac{p}{q} \sum_{k=-\infty}^{\infty} \alpha_k^q (x_{k+1} - x_k)^{\frac{q}{p}} \\ &= B^q \frac{p}{q} \sum_{k=-\infty}^{\infty} \alpha_k^q \left(\sum_{\{m: |f(2^m)|=\alpha_k\}} 2^m \right)^{\frac{q}{p}} \leq B^q \frac{p}{q} \sum_{k=-\infty}^{\infty} \sum_{\{m: |f(2^m)|=\alpha_k\}} |f(2^m)|^q 2^{\frac{mq}{p}} \\ &= B^q \frac{p}{q} \sum_{m \in \mathbb{Z}} |f(2^m)|^q 2^{\frac{mq}{p}} = 2B^q \frac{p}{q} \sum_{m \in \mathbb{Z}} \int_{2^{m-1}}^{2^m} |f(2^m)|^{\frac{q}{p}} 2^{m(\frac{q}{p}-1)} dx \\ &\leq 2B^{2q} \frac{p}{q} \int_0^\infty |f(x)|^q x^{\frac{q}{p}-1} dx. \square \end{aligned}$$

Theorem 4.10. Let $B > 1$, $f \in GM_1(B)$. If $0 < p < \infty$, $0 < q < \infty$ then:

$$\|f\|_{L(p,q)} \sim \left(\sum_{k=-\infty}^{\infty} 2^{\frac{kq}{p}} (f^*(2^k))^q \right)^{\frac{1}{q}} \sim \left(\sum_{k=-\infty}^{\infty} 2^{\frac{kq}{p}} |f(2^k)|^q \right)^{\frac{1}{q}} \sim \|f\|_{L_{w(p,q)}^q}. \quad (34)$$

If $0 < p \leq \infty$ then

$$\|f\|_{L(p,\infty)} \sim \sup_{k \in \mathbb{Z}} \left\{ 2^{\frac{k}{p}} |f(2^k)| \right\} \sim \sup_{k \in \mathbb{Z}} \left\{ 2^{\frac{k}{p}} |f^*(2^k)| \right\} \sim \|f\|_{L_{w(p,\infty)}^\infty}. \quad (35)$$

Proof.

First, show that $\|f\|_{L(p,q)} \sim \|f\|_{L_{w(p,q)}^q}$ for all $0 < q \leq \infty$.

The inequality $\|f\|_{L^q_{w(p,q)}} \leq B^2 C(p, q) \|f\|_{L(p,q)}$ follows from Lemma 4.8, more specifically, for $0 < p < \infty$, $0 < q < \infty$ it is a consequence of (31), and for $0 < p \leq \infty$, $q = \infty$ it is a consequence of (33).

The inequality $\|f\|_{L(p,q)} \leq B^2 C(p, q) \|f\|_{L^q_{w(p,q)}}$ follows from Theorem 3.4 for all $0 < p \leq q \leq \infty$ and it is a consequence of Lemma 4.9 for all $0 < q < p$.

$f^* \in GMS_1(1)$ as a decreasing function, and so by Lemma 4.8, if $0 < p < \infty$ then $\left(\sum_{k=-\infty}^{\infty} 2^{\frac{kq}{p}} |f^*(2^k)|^q \right)^{\frac{1}{q}} \sim \|f\|_{L(p,q)}$, and $\sup_{k \in \mathbb{Z}} \left\{ 2^{\frac{k}{p}} |f^*(2^k)| \right\} \sim \|f\|_{L(p,\infty)}$, that is, the remaining equivalences in (34) and (35) follow. $\square$

5. Properties of $GM_{\alpha,\varphi}$ and $GMS_{\alpha,\varphi}$.

Definition 5.1. For each $f \in L^1_{loc}(\mathbb{R}^+, \mu)$, where μ is a Borel measure, define for each x such that $\mu((0, x]) > 0$, the *average of f on $(0, x]$* , as

$$\sigma_x(f, \mu) := \frac{1}{\mu((0, x])} \int_{(0, x]} f d\mu. \quad (36)$$

Formula (36) defines a function of x , which we shall write as $\sigma(f, \mu)$. If f or μ are clear from the context, we may omit them from the notation.

Lemma 5.2. Let (Ω, Σ, μ) be a measure space, $-\infty < c < d < \infty$, and assume that $g : \Omega \times \mathbb{R} \rightarrow \mathbb{C}$ is such that for all $c \leq x \leq d$, $g(\cdot, x)$ is a μ -integrable function. Then:

$$V_{\int_{\Omega} g(\omega, \cdot) d\mu}([c, d]) \leq \int_{\Omega} V_{g(\omega, \cdot)}([c, d]) d\mu. \quad (37)$$

Lemma 5.3. Let (Ω, Σ, μ) be a measure space, and assume that for a.e. $\omega \in \Omega$, $g(\omega, \cdot) \in GM_{\alpha,\varphi}(B)$. Then

$$f(\cdot) := \int_{\Omega} g(\omega, \cdot) d\mu \in GM_{\alpha,\varphi}\left(\frac{B}{\cos \varphi}\right). \quad (38)$$

Corollary 5.3. $GM_{\alpha,\varphi}$ is a cone in GM . Moreover, if $f_j \in GM_{\alpha,\varphi}(B)$, $1 \leq j \leq N$ then:

$$\sum_{j=1}^N f_j \in GM_{\alpha,\varphi}\left(\frac{B}{\cos \varphi}\right). \quad (39)$$

Similarly, $GMS_{\alpha,\varphi}$ is a cone in GMS and for $b_n = \sum_{j=1}^N a_{n,j}$, where for each j the sequence $\{a_{n,j}\}_{n=1}^{\infty}$ is in $GMS(B)$, we have that

$$\{b_n\}_{n=1}^{\infty} \in GMS_{\alpha,\varphi}\left(\frac{B}{\cos \varphi}\right). \quad \square \quad (40)$$

Lemma 5.4. Assume that for a.e. $x > 0$, $f(x) \in S_{\alpha, \varphi}$, and that μ is a Borel measure on $\mathbb{R}^+$ for which there is $A \geq 0$ such that for all $a > A$, $\mu((0, a]) > 0$. Then, $\forall x > A$:

$$\sigma_x(|f|) \leq \frac{1}{\cos \varphi} |\sigma_x(f)|. \quad (41)$$

If, moreover, $f \in GM_1(B)$ and $\exists K > 0$ such that $\forall a > 0$,

$$\mu((0, a]) \leq K\mu((a, 2a]) \quad (42)$$

then, $\forall x > A$:

$$|f(x)| \leq \frac{(K+1)B}{\cos \varphi} |\sigma_x(f)|. \quad (43)$$

Proof.

(41) follows from Lemma 1.19:

$$\sigma_x(|f|) = \frac{1}{\mu((0, x])} \int_{(0, x]} |f| d\mu \leq \frac{1}{\mu((0, x]) \cos \varphi} \left| \int_{(0, x]} f d\mu \right| = \frac{1}{\cos \varphi} |\sigma_x(f)|.$$

Since $f \in GM_1(B)$, it follows that $|f(x)| \leq B|f(t)|$, $\forall t \in [\frac{x}{2}, x]$. Furthermore, by (42) for all $x > 0$:

$$\mu((0, x]) = \mu((0, \frac{x}{2}]) + \mu((\frac{x}{2}, x]) \leq (K+1)\mu((\frac{x}{2}, x]).$$

Therefore, for $x > A$:

$$\begin{aligned} |f(x)| &\leq \frac{B}{\mu((\frac{x}{2}, x])} \int_{(\frac{x}{2}, x]} |f(t)| d\mu \leq \frac{(K+1)B}{\mu((0, x])} \int_{(0, x]} |f(t)| d\mu \\ &= (K+1)B\sigma_x(|f|) \leq \frac{(K+1)B}{\cos \varphi} |\sigma_x(f)|. \square \end{aligned}$$

Analogously,

Lemma 5.5. Let $a_k \in S_{\alpha, \varphi}$, $\forall k \geq 1$. Then, $\forall n \geq 1$:

$$\sigma_n(\{|a_k|\}) \leq \frac{1}{\cos \varphi} |\sigma_n(\{a_k\})|.$$

If, moreover, $\{a_k\} \in GMS_1(B)$, then, $\forall n \geq 1$:

$$|a_n| \leq \frac{2B}{\cos \varphi} |\sigma_n(\{a_k\})|.$$

Lemma 5.6. Let $f \in GM_{\alpha, \varphi}(B)$ and assume that μ is a Borel measure on $\mathbb{R}^+$, for which there are $A \geq 0$, $K > 0$ such that for all $a > A$,

$$\begin{cases} \mu((0, a]) > 0 \\ \mu((a, 2a]) \leq K\mu((0, a]) \leq K^2\mu((0, a]) \end{cases}. \quad (44)$$

Then $\forall x > A$, $\sigma_x(f, \mu) \in S_{\alpha, \varphi}$. Moreover, there is $B' = B'(B, K, \varphi) > 0$ such that for all $x > A$, $\sigma(f, \mu)$ satisfies

$$V_{\sigma(f, \mu)}([x, 2x]) \leq B' |\sigma_x(f, \mu)|. \quad (45)$$

In particular, if $A = 0$ then $\sigma(f, \mu) \in GM_{\alpha, \varphi}(B')$.

Proof.

Clearly, $\sigma_x(f, \mu) \in S_{\alpha, \varphi}$.

Let $a > A$. Define, for $x \geq a$:

$$g(t, x) = f(t) \frac{I_{(0, x]}(t)}{\mu((0, x])}. \quad (46)$$

Since $f \in L^1_{loc}(\mathbb{R}^+, \mu)$, it follows that $g(\cdot, x) \in L^1(\mathbb{R}^+, \mu)$. Furthermore, $\sigma_x(f) = \int_{\mathbb{R}^+} g(t, x) d\mu$. By Lemma 5.2:

$$V_{\sigma(f)}([a, 2a]) = V_{\int_{\mathbb{R}^+} g(t, \cdot) d\mu}([a, 2a]) \leq \int_{\mathbb{R}^+} V_{g(t, \cdot)}([a, 2a]) d\mu.$$

Clearly, $g(t, x) = f(t) \frac{I_{[t, \infty)}(x)}{\mu((0, x])}$ is a decreasing function of x on $[t, \infty)$. Consider the following three cases: $0 \leq t \leq a$, $a < t \leq 2a$, $t > 2a$.

If $0 \leq t \leq a$ then

$$V_{g(t, x)}([a, 2a]) = V_{\frac{f(t)}{\mu((0, x])}}([a, 2a]) = \frac{|f(t)|}{\mu((0, a])} - \frac{|f(t)|}{\mu((0, 2a])}.$$

If $a < t \leq 2a$, observe that for $a \leq x < t$, $g(t, x) = 0$, and $g(t, t) = \frac{|f(t)|}{\mu((0, t])}$, therefore, $V_{g(t, x)}([a, t]) = \frac{|f(t)|}{\mu((0, t])}$, and so:

$$\begin{aligned} V_{g(t, x)}([a, 2a]) &= V_{g(t, x)}([a, t]) + V_{g(t, x)}([t, 2a]) \\ &= \frac{|f(t)|}{\mu((0, t])} + V_{\frac{f(t)}{\mu((0, x])}}([t, 2a]) \\ &= \frac{|f(t)|}{\mu((0, a])} + \left(\frac{|f(t)|}{\mu((0, t])} - \frac{|f(t)|}{\mu((0, 2a])} \right) = \frac{2|f(t)|}{\mu((0, t])} - \frac{|f(t)|}{\mu((0, 2a])}. \end{aligned}$$

Finally, if $t > 2a$ then $g(t, x) = 0$, $\forall x \in [a, 2a]$, and so, $V_{g(t, x)}([a, 2a]) = 0$. Therefore:

$$\begin{aligned} &\int_{\mathbb{R}^+} V_{g(t, \cdot)}([a, 2a]) d\mu = \\ &= \int_{(0, a]} \left(\frac{1}{\mu((0, a])} - \frac{1}{\mu((0, 2a])} \right) |f(t)| d\mu \end{aligned}$$

$$\begin{aligned}
& + \int_{(a, 2a]} \left(\frac{2}{\mu((0, t])} - \frac{1}{\mu((0, 2a])} \right) |f(t)| d\mu \\
& \leq \left(\frac{2}{\mu((0, a])} - \frac{1}{\mu((0, 2a])} \right) \int_{(0, 2a]} |f(t)| d\mu \\
& = \frac{2\mu((a, 2a]) + \mu((0, a])}{\mu((0, a])} \sigma_{2a}(|f|) \leq (2K + 1) \sigma_{2a}(|f|).
\end{aligned}$$

Observe that by Lemma 1.22 for $f \in GM(B)$ and $a < t \leq 2a$, $|f(t)| \leq 2B|f(a)|$, and so:

$$\begin{aligned}
\sigma_{2a}(|f|) &= \frac{1}{\mu((0, 2a])} \left(\int_{(0, a]} |f| d\mu + \int_{(a, 2a]} |f| d\mu \right) \\
&\leq \frac{1}{\mu((0, a])} \int_{(0, a]} |f| d\mu + \frac{1}{\mu((a, 2a])} \int_{(a, 2a]} 2B|f(a)| d\mu = \sigma_a(|f|) + 2B|f(a)|.
\end{aligned}$$

Since μ satisfies (42) and $f \in GM_{\alpha, \varphi}$, it follows from Lemma 5.4 that:

$$\sigma_a(|f|) \leq \frac{1}{\cos \varphi} |\sigma_a(f)| \quad \text{and} \quad |f(a)| \leq \frac{(K+1)B}{\cos \varphi} |\sigma_a(f)|,$$

and so, $\sigma_{2a}(|f|) \leq \frac{1 + 2(K+1)B^2}{\cos \varphi} |\sigma_a(f)|$. Therefore,

$$V_{\sigma(f, \mu)}([a, 2a]) \leq (2K+1) \sigma_{2a}(|f|) \leq \frac{(2K+1)(1 + 2(K+1)B^2)}{\cos \varphi} |\sigma_a(f, \mu)|. \square$$

6. Fourier series with general monotone coefficients.

Theorem 6.1. Assume that $a_k \in \mathbb{C}$, $1 \leq m \leq N$. Then, $\forall x \in (0, \pi]$:

$$\left| \sum_{k=m}^N a_k e^{ikx} \right| \leq \frac{4\pi}{x} \left(\frac{|a_m|}{2} + \sum_{k=m}^{N-1} |a_{k+1} - a_k| \right). \quad (47)$$

Proof.

Since for $x \in (0, \pi]$,

$$\left| \sum_{k=0}^N e^{ikx} \right| \leq \frac{1}{\sin \frac{x}{2}} \leq \frac{\pi}{x}, \quad (48)$$

it follows, using summation by parts,

$$\sum_{k=m}^N a_k e^{ikx} = a_N \sum_{k=m}^N e^{ikx} - \sum_{k=m}^{N-1} (a_{k+1} - a_k) \sum_{j=m}^k e^{ijx},$$

and so

$$\left| \sum_{k=m}^N a_k e^{ikx} \right| \leq \frac{2\pi}{x} \left(|a_N| + \sum_{k=m}^{N-1} |a_{k+1} - a_k| \right).$$

Since $a_N = a_m + \sum_{k=m}^{N-1} (a_{k+1} - a_k)$, it follows that

$$\left| \sum_{k=m}^N a_k e^{ikx} \right| \leq \frac{2\pi}{x} \left(|a_m| + 2 \cdot \sum_{k=m}^{N-1} |a_{k+1} - a_k| \right). \quad (49)$$

Corollary 6.1. Assume $\{a_k\} \in GMS_2(B)$. Then $\forall x \in (0, \pi]$, $\forall 1 \leq m \leq N$,

$$\left| \sum_{k=m}^N a_k e^{ikx} \right| \leq \frac{6\pi B}{x} \left(|a_m| + \sum_{k=m+1}^N \frac{|a_k|}{k} \right). \quad (50)$$

Theorem 6.2. Assume $\{c_k\} \in GMS_2(B)$. If $f(x) = \sum_{k=1}^{\infty} c_k e^{ikx}$, then:

$$\|f\|_{L^1(0, \pi)} \leq 2\pi |c_1| + 27\pi B \|\{c_n\}\|_{l^1_{\frac{\ln k}{k}}}. \quad (51)$$

Proof.

For each $N \geq 2$, denote $f_N(x) = \sum_{k=1}^N c_k e^{ikx}$. Then:

$$\begin{aligned} \|f_N\|_{L^1(0, \pi)} &= \sum_{n=1}^{\infty} \int_{\frac{\pi}{n+1}}^{\frac{\pi}{n}} \left| \sum_{k=1}^N c_k e^{ikx} \right| dx \\ &\leq \sum_{n=1}^{N-1} \int_{\frac{\pi}{n+1}}^{\frac{\pi}{n}} \left| \sum_{k=1}^N c_k e^{ikx} \right| dx + \sum_{n=N}^{\infty} \int_{\frac{\pi}{n+1}}^{\frac{\pi}{n}} \left(\sum_{k=1}^N |c_k| \right) dx \\ &\leq \sum_{n=1}^{N-1} \int_{\frac{\pi}{n+1}}^{\frac{\pi}{n}} \left(\sum_{k=1}^n |c_k| + \left| \sum_{k=n+1}^N c_k e^{ikx} \right| \right) dx + \frac{\pi}{N} \sum_{k=1}^N |c_k|. \end{aligned}$$

Applying (50) for $n < N-1$:

$$\left| \sum_{k=n+1}^N c_k e^{ikx} \right| \leq \frac{6\pi B}{x} \left(|c_{n+1}| + \sum_{k=n+2}^N \frac{|c_k|}{k} \right).$$

If $n = N-1$ then $\left| \sum_{k=n+1}^N c_k e^{ikx} \right| = |c_N|$, and so, for $n \leq N-1$,

$$\left| \sum_{k=n+1}^N c_k e^{ikx} \right| \leq \frac{6\pi B}{x} \left(|c_{n+1}| + \sum_{k=n+1}^N \frac{|c_k|}{k} \right).$$

Therefore:

$$\begin{aligned}\|f_N\|_{L^1(0,\pi)} &\leq \sum_{n=1}^{N-1} \int_{\frac{\pi}{n+1}}^{\frac{\pi}{n}} \left[\sum_{k=1}^n |c_k| + \frac{6\pi B}{x} \left(|c_{n+1}| + \sum_{k=n+1}^N \frac{|c_k|}{k} \right) \right] dx + \frac{\pi}{N} \sum_{k=1}^N |c_k| \\ &= I_1 + I_2 + I_3 + \frac{\pi}{N} \sum_{k=1}^N |c_k|,\end{aligned}$$

where, using summation by parts:

$$\begin{aligned}I_1 &= \sum_{n=1}^{N-1} \int_{\frac{\pi}{n+1}}^{\frac{\pi}{n}} \left(\sum_{k=1}^n |c_k| \right) dx = \sum_{n=1}^{N-1} \left(\frac{\pi}{n} - \frac{\pi}{n+1} \right) \sum_{k=1}^n |c_k| \\ &= -\frac{\pi}{N} \sum_{n=1}^N |c_n| + \pi \sum_{n=1}^N \frac{|c_n|}{n} \leq \pi \sum_{n=1}^N \frac{|c_n|}{n}.\end{aligned}$$

It follows that

$$\begin{aligned}I_1 &\leq \pi \left(|c_1| + \sum_{n=2}^N \frac{|c_n|}{n} \right) \leq \pi |c_1| + \frac{\pi}{\ln 2} \sum_{n=2}^N |c_n| \frac{\ln n}{n}. \\ I_2 &= \sum_{n=1}^{N-1} \int_{\frac{\pi}{n+1}}^{\frac{\pi}{n}} \frac{6\pi B}{x} |c_{n+1}| dx = \\ &= 6\pi B \sum_{n=1}^{N-1} |c_{n+1}| \ln \frac{n+1}{n} \leq 6\pi B \sum_{n=1}^{N-1} \frac{|c_{n+1}|}{n} = 6\pi B \sum_{n=2}^N \frac{|c_n|}{n-1} \\ &\leq 12\pi B \sum_{n=2}^N \frac{|c_n|}{n} \leq \frac{12\pi B}{\ln 2} \sum_{n=2}^N |c_n| \frac{\ln n}{n}; \\ I_3 &= \sum_{n=1}^{N-1} \int_{\frac{\pi}{n+1}}^{\frac{\pi}{n}} \frac{6\pi B}{x} \left(\sum_{k=n+1}^N \frac{|c_k|}{k} \right) dx = 6\pi B \sum_{n=1}^{N-1} \left(\sum_{k=n+1}^N \frac{|c_k|}{k} \right) \int_{\frac{\pi}{n+1}}^{\frac{\pi}{n}} \frac{dx}{x} \\ &= 6\pi B \sum_{n=1}^{N-1} \sum_{k=n}^{N-1} \frac{|c_{k+1}|}{k+1} \ln \frac{n+1}{n} = 6\pi B \sum_{k=1}^{N-1} \frac{|c_{k+1}|}{k+1} \sum_{n=1}^k \ln \frac{n+1}{n} \\ &= 6\pi B \sum_{k=1}^{N-1} \frac{|c_{k+1}|}{k+1} \ln(k+1) = 6\pi B \sum_{k=2}^N |c_k| \frac{\ln k}{k}.\end{aligned}$$

Combining the estimates for I_j and taking into account that

$$\frac{\pi}{N} \sum_{k=1}^N |c_k| \leq \pi |c_1| + \frac{\pi}{N \ln 2} \sum_{k=2}^N |c_k| \ln k \leq \pi |c_1| + \frac{\pi B}{\ln 2} \sum_{k=2}^N |c_k| \frac{\ln k}{k},$$

we have:

$$\|f_N\|_{L^1(0,\pi)} \leq 2\pi |c_1| + 27\pi B \sum_{k=2}^N |c_k| \frac{\ln k}{k}. \square$$

Theorem 6.3. Assume $\{c_k\} \in GMS_2(B)$. If $f(x) = \sum_{k=1}^{\infty} c_k e^{ikx}$, then

$$\|f\|_{L(1,\infty)(0,\pi)} \leq 6\pi B \|\{c_n\}\|_{l_{1/k}^1}. \quad (52)$$

Proof.

Let $N \geq 1$. Denote $f_N(x) = \sum_{k=1}^N c_k e^{ikx}$. Then, by (50),

$$|f_N(x)| = \left| \sum_{k=1}^N c_k e^{ikx} \right| \leq \frac{6\pi B}{x} \left(|c_1| + \sum_{k=2}^N \frac{|c_k|}{k} \right) = \frac{6\pi B}{x} \sum_{k=1}^N \frac{|c_k|}{k}.$$

Therefore, for $\forall \alpha > 0$:

$$\begin{aligned} \lambda(\{x : |f_N(x)| > \alpha\}) &\leq \lambda\left(\left\{x : \frac{6\pi B}{x} \sum_{k=1}^N \frac{|c_k|}{k} > \alpha\right\}\right) \\ &\leq \lambda\left(\left\{x : \frac{6\pi B}{x} \|\{c_k\}\|_{l_{1/k}^1} > \alpha\right\}\right) = \frac{6\pi B}{\alpha} \|\{c_k\}\|_{l_{1/k}^1}, \end{aligned}$$

proving (52) $\square$

The result of Theorem 6.3 can be interpreted as follows. Let T be the operator mapping GMS_2 to functions over $\mathbb{R}$, defined as

$$T(\{c_n\})(x) := \sum_{k=1}^{\infty} c_k e^{ikx}. \quad (53)$$

Then

$$T : l_{\frac{1}{k}}^1 \cap GMS_2 \rightarrow L(1, \infty). \quad (54)$$

Theorem 6.4. Let $1 < p < \infty$, $0 < q \leq \infty$. If c_k are trigonometric Fourier coefficients of f and $\{c_k\} \in GMS_{\alpha,\varphi}(B)$ then $\exists C = C(p, q, \varphi)$ such that

$$\|\{c_n\}\|_{l(p',q)} \leq C \|f\|_{L(p,q)(0,2\pi)}. \quad (55)$$

Proof.

By Lemma 5.5, $|c_n| \leq \frac{2B}{\cos \varphi} |\sigma_n|$, and so $\|\{c_n\}\|_{l(p',q)} \leq \frac{2B}{\cos \varphi} \|\sigma_n\|_{l(p',q)}$. By (28),

$$\|\{c_n\}\|_{l(p',q)} \leq \frac{C'(p,q)}{\cos \varphi} \|f\|_{L(p,q)(0,2\pi)}. \square$$

7. Interpolation in $GMS_{\alpha,\varphi}$.

The following Lemma is a special case of a result of Yu. Brudnui and N. Krugljak ([4], Corollary 3.1.26).

Lemma 7.1.

$$K\left(t, \{c_n\}, l^{\frac{1}{k}}, l^1\right) \sim t \sum_{n \leq \frac{1}{t}} |c_n| + \sum_{n > \frac{1}{t}} \frac{|c_n|}{n} \quad (56)$$

Lemma 7.2. Let $\{a_n\}, \{c_n\} \in GMS(B)$. Assume that $N \geq 1$ and for some $\gamma > 0$, $|c_N| \leq \gamma |a_N|$. Define

$$b_n = \begin{cases} a_n & \text{if } n \leq N \\ c_n & \text{if } n > N \end{cases}.$$

Then $\{b_n\} \in GMS(3B + 6B^2\gamma)$.

Proof.

If $n = N = 1$ then

$$\sum_{k=n}^{2n-1} |b_{k+1} - b_k| = |a_1 - c_2| = |a_N - c_{N+1}|.$$

If $n = N > 1$ then

$$\sum_{k=n}^{2n-1} |b_{k+1} - b_k| = |a_N - c_{N+1}| + \sum_{k=N+1}^{2N-1} |c_{k+1} - c_k|.$$

If $n = \frac{N+1}{2} > 1$ then

$$\sum_{k=n}^{2n-1} |b_{k+1} - b_k| = \sum_{k=n}^{N-1} |a_{k+1} - a_k| + |a_N - c_{N+1}|.$$

If $\frac{N+1}{2} < n < N$ then

$$\sum_{k=n}^{2n-1} |b_{k+1} - b_k| = \sum_{k=n}^{N-1} |a_{k+1} - a_k| + |a_N - c_{N+1}| + \sum_{k=N+1}^{2n-1} |c_{k+1} - c_k|.$$

In all four cases above:

$$\sum_{k=n}^{2n-1} |b_{k+1} - b_k| \leq \sum_{k=n}^{2n-1} |a_{k+1} - a_k| + |a_N - c_{N+1}| + \sum_{k=N}^{2N-1} |c_{k+1} - c_k|,$$

and $\frac{N}{2} < n \leq N$. By (19), $|a_N| \leq 2B|a_n|$ and $|c_{N+1}| \leq 2B|c_N|$, and so:

$$\begin{aligned} \sum_{k=n}^{2n-1} |b_{k+1} - b_k| &\leq \sum_{k=n}^{2n-1} |a_{k+1} - a_k| + |a_N| + |c_{N+1}| + \sum_{k=N}^{2N-1} |c_{k+1} - c_k| \\ &\leq B|a_n| + 2B|a_n| + (2B + B)|c_N| \leq 3B|a_n| + 3B\gamma|a_n| \\ &\leq 3B|a_n| + 6B^2\gamma|a_n| = (3B + 6B^2\gamma)|b_n|. \end{aligned}$$

Two cases remain. If $n < \frac{N+1}{2}$ then $2n-1 \leq N-1$, and so:

$$\sum_{k=n}^{2n-1} |b_{k+1} - b_k| = \sum_{k=n}^{2n-1} |a_{k+1} - a_k| \leq B|a_n| = B|b_n|.$$

If $n > N$ then

$$\sum_{k=n}^{2n-1} |b_{k+1} - b_k| = \sum_{k=n}^{2n-1} |c_{k+1} - c_k| \leq B|c_n| = B|b_n|. \square$$

Lemmas 7.1 and 7.2 imply:

Theorem 7.3.

$$K\left(t, \{c_n\}, l_{\frac{1}{k}}^1 \cap GMS_{\alpha, \varphi}, l^1\right) \leq \frac{9}{2} K\left(t, \{c_n\}, l_{\frac{1}{k}}^1, l^1\right). \quad (57)$$

Proof.

Let $\{c_n\} \in GMS_{\alpha, \varphi}(B) \cap \left(l_{\frac{1}{k}}^1 + l^1\right) = GMS_{\alpha, \varphi}(B) \cap l_{\frac{1}{k}}^1$.

1) If $t > 1$ then, by (56), $K\left(t, \{c_n\}, l_{\frac{1}{k}}^1, l^1\right) = \sum_{n=1}^{\infty} \frac{|c_n|}{n} = \|\{c_n\}\|_{l_{\frac{1}{k}}^1}$, and (57)

follows.

2) Assume that $t \leq 1$.

Let $N = 1 + \left\lfloor \frac{1}{t} \right\rfloor$, $\sigma_N = \frac{1}{N} \sum_{k=1}^N |c_k|$, and define $a_n = \frac{n}{N} \sigma_N e^{i\alpha}$ for all

$n \geq 1$. Then $\sum_{k=n}^{2n-1} |a_k - a_{k+1}| = |a_{2n} - a_n| = \frac{n}{N} \sigma_N = |a_n|$, and so $\{a_n\} \in GMS(1)$. By Corollary 5.3, $\{a_n\} \in GMS_{\alpha, \varphi}(1)$. Observe that for $2 \leq N \leq 4$,

$\#\{k \in \mathbb{Z} : \frac{N}{4} < k \leq \frac{N}{2}\} = 1 > \frac{N}{5}$. Assume that $N > 4$ and let $\theta = \frac{N}{4} - \lfloor \frac{N}{4} \rfloor$. If $\theta < \frac{1}{2}$ then $\theta \leq \frac{1}{4}$, and so:

$$\#\left\{k \in \mathbb{Z} : \frac{N}{4} < k \leq \frac{N}{2}\right\} = \left\lfloor \frac{N}{4} \right\rfloor \geq \frac{N}{4} - \frac{1}{4} \geq \frac{N}{4} - \frac{N/5}{4} = \frac{N}{5}.$$

If $\theta \geq \frac{1}{2}$ then

$$\#\left\{k \in \mathbb{Z} : \frac{N}{4} < k \leq \frac{N}{2}\right\} \geq \left\lfloor \frac{N}{4} \right\rfloor + 1 > \frac{N}{4} > \frac{N}{5}.$$

Applying (11) to $\{c_n\}$, for all $n \in \left[\frac{N}{2}, N\right]$ we have:

$$\begin{aligned} |a_n| &= \frac{n}{N} \sigma_N \geq \frac{1}{2} \sigma_N \geq \frac{1}{2} \cdot \frac{1}{N} \sum_{k \in (\frac{N}{4}, \frac{N}{2}]} |c_k| \geq \frac{1}{2} \cdot \frac{1}{N} \sum_{k \in (\frac{N}{4}, \frac{N}{2}]} \frac{1}{B} \left| c_{\lceil \frac{N}{2} \rceil} \right| \\ &\geq \frac{1}{2N} \cdot \frac{N}{5} \cdot \frac{1}{B} \left| c_{\lceil \frac{N}{2} \rceil} \right| = \frac{1}{10B} \left| c_{\lceil \frac{N}{2} \rceil} \right| \geq \frac{1}{10B^2} |c_n|, \end{aligned}$$

in particular, $|c_N| \leq 10B^2 |a_N|$.

Define

$$b_n = \begin{cases} a_n & \text{if } n \leq N \\ c_n & \text{if } n > N \end{cases} \quad ; \quad d_n = \begin{cases} c_n - a_n & \text{if } n \leq N \\ 0 & \text{if } n > N \end{cases}.$$

By Lemma 7.2, $\{b_n\} \in GMS_{\alpha, \varphi}(3B + 6B^2 \cdot 10B^2) \subset GMS_{\alpha, \varphi}(63B^4)$ and $b_n + d_n = c_n$, all n . Furthermore:

$$\begin{aligned} \|\{b_n\}\|_{l^1_{\frac{1}{N}}} &= \left(\sum_{n=1}^N + \sum_{n=N+1}^{\infty} \right) \frac{|b_n|}{n} \\ &= \sum_{n=1}^N \frac{|a_n|}{n} + \sum_{n=N+1}^{\infty} \frac{|c_n|}{n} = \sigma_N + \sum_{n=N+1}^{\infty} \frac{|c_n|}{n}; \\ \|\{d_n\}\|_{l^1} &= \sum_{n=1}^N |c_n - a_n| \\ &\leq \sum_{n=1}^N |a_n| + \sum_{n=1}^N |c_n| = \frac{\sigma_N}{N} \sum_{n=1}^N n + \sum_{n=1}^N |c_n| \\ &= \sigma_N \frac{N+1}{2} + N\sigma_N = \sigma_N \frac{3N+1}{2}. \end{aligned}$$

Therefore, observing that $t \leq \frac{1}{N-1} \leq \frac{2}{N}$, we obtain:

$$\|\{b_n\}\|_{l^1_{\frac{1}{N}}} + t \|\{d_n\}\|_{l^1}$$

$$\begin{aligned}
&= \sigma_N \left(1 + \frac{3N+1}{2} t \right) + \sum_{n=N+1}^{\infty} \frac{|c_n|}{n} \leq \sigma_N \left(1 + \frac{3N+1}{N} \right) + \sum_{n=N+1}^{\infty} \frac{|c_n|}{n} \\
&\leq \frac{9}{2} \sigma_N + \sum_{n=N+1}^{\infty} \frac{|c_n|}{n} = \frac{9}{2N} \sum_{n=1}^N |c_n| + \sum_{n=N+1}^{\infty} \frac{|c_n|}{n} \\
&= \frac{9}{2N} \left(\sum_{n=1}^{\lfloor \frac{1}{t} \rfloor} |c_n| + |c_N| \right) + \sum_{n=N+1}^{\infty} \frac{|c_n|}{n} \\
&\leq \frac{9}{2} \left(t \sum_{n=1}^{\lfloor \frac{1}{t} \rfloor} |c_n| + \sum_{n=N}^{\infty} \frac{|c_n|}{n} \right) = \frac{9}{2} K \left(t, \{c_n\}, l_{\frac{1}{k}}^1, l^1 \right),
\end{aligned}$$

in the last line we applied $N = \lfloor \frac{1}{t} \rfloor + 1 \in (\frac{1}{t}, \frac{1}{t} + 1]$. $\square$

B. Booton [3] gave a different proof of the following theorem for $1 < p, q \geq 1$ and $\{c_k\} \in GMS^+$.

Theorem 7.4. Let $1 < p < \infty$, $0 < q \leq \infty$. Assume $\{c_k\} \in GMS_{\alpha, \varphi}$. Let $f(x) = \sum_{k=1}^{\infty} c_k e^{ikx}$. Then $\exists C = C(p, q)$ such that

$$\|f\|_{L(p, q)(0, \pi)} \leq C \|\{c_n\}\|_{l(p', q)}. \quad (58)$$

Proof.

Let $X = l_{\frac{1}{k}}^1$ and let w be the weight: $w(k) = k$. Then $X_w = l^1$. Both X and X_w can be considered as quasi-normed monoids, and so, by Gilbert's theorem (2.6),

$$\begin{aligned}
&\|\{c_k\}\|_{(X, X_w)_{\theta, q; K}} \sim \left(\int_0^{\infty} (t^{-\theta} \|\{c_k\} \cdot \sigma_t(k)\|_X)^q \frac{dt}{t} \right)^{1/q} \\
&= \left(\int_0^{\infty} \left(t^{-\theta} \sum_{k=1}^{\infty} tk \sigma(tk) \frac{|c_k|}{k} \right)^q \frac{dt}{t} \right)^{1/q} = \left(\int_0^{\infty} \left(t^{1-\theta} \sum_{k=1}^{\infty} \sigma(tk) |c_k| \right)^q \frac{dt}{t} \right)^{1/q} \\
&= \left(\int_0^{\infty} \left(t^{\theta-1} \sum_{k=1}^{\infty} \sigma\left(\frac{k}{t}\right) |c_k| \right)^q \frac{dt}{t} \right)^{1/q}.
\end{aligned}$$

Since $\sigma = I_{[1, 2]}$ satisfies the conditions of Gilbert's theorem,

$$\left(\int_0^{\infty} \left(t^{\theta-1} \sum_{k=1}^{\infty} \sigma\left(\frac{k}{t}\right) |c_k| \right)^q \frac{dt}{t} \right)^{1/q} \sim \left(\int_0^{\infty} \left(t^{\theta-1} \sum_{t \leq k < 2t} |c_k| \right)^q \frac{dt}{t} \right)^{1/q}.$$

$\{c_k\} \in GMS_1(B)$ for some $B \geq 1$, and so:

$$\sum_{t \leq k < 2t} |c_k| \leq \#\{k : t \leq k < 2t\} \cdot B |c_{\lceil t \rceil}| \leq 2Bt |c_{\lceil t \rceil}|.$$

Therefore, for $q < \infty$:

$$\begin{aligned} & \left(\int_0^\infty \left(t^{\theta-1} \sum_{t \leq k < 2t} |c_k| \right)^q \frac{dt}{t} \right)^{1/q} \leq \left(\int_0^\infty (2Bt^\theta |c_{\lceil t \rceil}|)^q \frac{dt}{t} \right)^{1/q} \\ & = 2B \left(\sum_{m=1}^\infty \int_{m-1}^m (t^\theta |c_{\lceil t \rceil}|)^q \frac{dt}{t} \right)^{1/q} = 2B \left(\sum_{m=1}^\infty |c_m|^q \int_{m-1}^m t^{\theta q} \frac{dt}{t} \right)^{1/q} \\ & \leq BC(\theta, q) \left(\sum_{m=1}^\infty |c_m|^q m^{\theta q - 1} \right)^{1/q} = BC(\theta, q) \|\{c_k\}\|_{l^q_w(\frac{1}{\theta}, q)}, \end{aligned}$$

where $C(\theta, q) = \max \left\{ 1, \frac{1}{\theta q}, 2^{1-\theta q} \right\}$.

Again, since $\{c_k\} \in GMS_1(B)$, it follows that for $t \leq k < 2t$, $|c_k| \geq \frac{1}{B} |c_{\lfloor 2t \rfloor}|$. Also:

If $t \leq \frac{1}{2}$ and $k \in \mathbb{N}$ then $\sigma\left(\frac{k}{t}\right) = 0$.

If $n - \frac{1}{2} < t \leq n$, where $n \in \mathbb{N}$, then

$$\#\{k : t \leq k < 2t\} = \#\{k : n \leq k \leq 2n - 1\} = n > t.$$

If $n < t \leq n + \frac{1}{2}$, where $n \in \mathbb{N}$, then

$$\#\{k : t \leq k < 2t\} = \#\{k : n + 1 \leq k \leq 2n\} = n > t - \frac{1}{2} > t - \frac{t}{2} = \frac{t}{2}.$$

Therefore, for $t \geq \frac{1}{2}$, $\sum_{t \leq k < 2t} |c_k| \geq \frac{t}{2B} |c_{\lfloor 2t \rfloor}|$, and so:

$$\begin{aligned} & \left(\int_0^\infty \left(t^{\theta-1} \sum_{t \leq k < 2t} |c_k| \right)^q \frac{dt}{t} \right)^{1/q} = \left(\int_{\frac{1}{2}}^\infty \left(t^{\theta-1} \sum_{t \leq k < 2t} |c_k| \right)^q \frac{dt}{t} \right)^{1/q} \\ & \geq \left(\int_{\frac{1}{2}}^\infty \left(\frac{t^\theta}{2B} |c_{\lfloor 2t \rfloor}| \right)^q \frac{dt}{t} \right)^{1/q} = \frac{1}{2B} \left(\sum_{m=1}^\infty \int_{\frac{m}{2}}^{\frac{m+1}{2}} (t^\theta |c_{\lfloor 2t \rfloor}|)^q \frac{dt}{t} \right)^{1/q} \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2B} \left(\sum_{m=1}^{\infty} |c_m|^q \int_{\frac{m}{2}}^{\frac{m+1}{2}} t^{\theta q} \frac{dt}{t} \right)^{1/q} \geq \frac{C(\theta, q)}{B} \left(\sum_{m=1}^{\infty} |c_m|^q m^{\theta q - 1} \right)^{1/q} \\
&= \frac{C(\theta, q)}{B} \|\{c_k\}\|_{l^q_{w(\frac{1}{\theta}, q)}}^q,
\end{aligned}$$

where $C(\theta, q) = \min \left\{ \frac{1}{2}, \left(\frac{1}{2} \right)^{\theta q - 1} \right\}$.

Therefore:

$$\|\{c_k\}\|_{(X, X_w)_{\theta, q; K}} \sim \|\{c_k\}\|_{l^q_{w(\frac{1}{\theta}, q)}}. \quad (59)$$

On the other hand, Theorem 7.3 shows that

$$\|\{c_k\}\|_{(X, X_w)_{\theta, q; K}} = \|\{c_k\}\|_{\left(l^1_{\frac{1}{K}}, l^1\right)_{\theta, q; K}} \sim \|\{c_k\}\|_{\left(l^1_{\frac{1}{K}} \cap GMS_{\alpha, \varphi}, l^1\right)_{\theta, q; K}}.$$

$l^1_{\frac{1}{K}} \cap GMS_{\alpha, \varphi}$ is a quasi-normed monoid in the vector space of sequences (see Corollary 5.3), with the norm $\|\cdot\|_{l^1_{\frac{1}{K}}}$. Assume that T is defined as in (53). By Theorem 6.3, T maps G_0 into $L(1, \infty)$, and $\|T\{c_n\}\|_{L(1, \infty)} \leq 6\pi B \|\{c_n\}\|_{l^1_{\frac{1}{K}}}$. Also, since a function with an absolutely convergent Fourier series is bounded, it follows that $T : G_1 \rightarrow L(\infty, \infty)$. Therefore,

$$\|T\{c_n\}\|_{(L(1, \infty), L(\infty, \infty))_{\theta, q; K}} \leq C(B) \|\{c_n\}\|_{(G_0, G_1)_{\theta, q; K}} \sim \|\{c_n\}\|_{(X, X_w)_{\theta, q; K}}.$$

Finally, by the interpolation theorem, $\forall \theta \in (0, 1), \forall q \in (0, \infty)$:

$$(L(1, \infty), L(\infty, \infty))_{\theta, q} = L(p, q),$$

where $p = \frac{1}{1 - \theta}$. That is to say, $\theta = \frac{1 - p}{p} = \frac{1}{p'}$.

Since $\{c_k\} \in GMS$ the conditions of Lemma 4.4 hold and, therefore, $|c_n| \leq Bc^*_{\lfloor \frac{n}{2} \rfloor}$. This implies that $\|\{c_k\}\|_{l^q_{w(p', q)}} \leq B'(p, q) \|\{c_k^*\}\|_{l^q_{w(p', q)}} = B'(p, q) \|\{c_k\}\|_{l(p, q)}$, thus obtaining (58). $\square$

Combining the results of Theorems 7.4 and 6.4, we obtain the main result of this section.

Theorem 7.5. Let $1 < p < \infty$, $0 < q \leq \infty$. Assume $\{c_k\} \in GMS_{\alpha, \varphi}$. Define $f(x) = \sum_{k=1}^{\infty} c_k e^{ikx}$. Then:

$$\|\{c_n\}\|_{l(p', q)} \sim \|f\|_{L(p, q)(0, \pi)}. \quad (60)$$

8. On Hardy's inequality for general monotone functions.

Lemma 8.1. If $f, g \in GM$ then $f \cdot g \in GM$.

Proof.

$$\begin{aligned}
V_{fg}([x, 2x]) &= \sup_{\{x_j\} \in \mathcal{P}([x, 2x])} \left\{ \sum_{j=0}^{n-1} |f(x_{j+1})g(x_{j+1}) - f(x_j)g(x_j)| \right\} \\
&\leq \sup_{\{x_j\} \in \mathcal{P}([x, 2x])} \left\{ \sum_{j=0}^{n-1} |f(x_{j+1})| |g(x_{j+1}) - g(x_j)| \right\} \\
&\quad + \sup_{\{x_j\} \in \mathcal{P}([x, 2x])} \left\{ \sum_{j=0}^{n-1} |g(x_j)| |f(x_{j+1}) - f(x_j)| \right\} \\
&\leq \sup_{[x, 2x]} |f(t)| \cdot V_g([x, 2x]) + \sup_{[x, 2x]} |g(t)| \cdot V_f([x, 2x]).
\end{aligned}$$

Since $f, g \in GM$ then $\exists B_1 > 0, B_2 > 0$ such that $\forall x > 0$:

$$\begin{cases} V_f([x, 2x]) \leq B_1 |f(x)| \\ V_g([x, 2x]) \leq B_2 |g(x)| \end{cases}.$$

Also, by (17), for all $x > 0$ and $t \in [x, 2x]$:

$$\begin{cases} |f(t)| \leq 2B_1 |f(x)| \\ |g(t)| \leq 2B_2 |g(x)| \end{cases}.$$

Therefore, $\forall x > 0$:

$$V_{fg}([x, 2x]) \leq 4B_1 B_2 |f(x)g(x)|,$$

that is, $fg \in GM(4B_1 B_2)$. $\square$

We use interpolation to give a very short proof of Hardy's inequality for general monotone functions, for $\forall \alpha > 0, \forall q > 0$.

Theorem 8.2 (Hardy's inequality for general monotone functions).

Let $\alpha > 0, 0 < q \leq \infty$, and $f \in GM^+$. Then

$$\left(\int_0^\infty \left(x^{-\alpha} \int_0^x f(t) \frac{dt}{t} \right)^q \frac{dx}{x} \right)^{\frac{1}{q}} \leq C(\alpha, q) \left(\int_0^\infty (x^{-\alpha} f(x))^q \frac{dx}{x} \right)^{\frac{1}{q}}. \quad (61)$$

Proof.

1) Consider $\alpha \leq 1$. Take $g(t) = \frac{f(t)}{t}$. By Lemma 8.1, since $\frac{1}{t} \in GM^+$, $g \in GM^+$.

Let T be defined for integrable on $[0, a]$ functions, for all $a > 0$, as

$$(Th)(x) = \frac{1}{x} \int_0^x h(t) dt.$$

Then $T : L^\infty \rightarrow L^\infty$ and $T : L(1, 1) \rightarrow L(1, \infty)$. Applying the interpolation theorem, we obtain

$$T : L(p, q) \rightarrow L(p, q),$$

where $1 < p < \infty$, $0 < q \leq \infty$.

This implies

$$\begin{aligned} \|Tg\|_{L(p,q)} &= \left(\int_0^\infty x^{\frac{q}{p}} \left(\left(\frac{1}{x} \int_0^x g(t) dt \right)^* \right)^q \frac{dx}{x} \right)^{\frac{1}{q}} \\ &\leq C(p, q) \left(\int_0^\infty x^{\frac{q}{p}} (g^*(x))^q \frac{dx}{x} \right)^{\frac{1}{q}} = C(p, q) \|g\|_{L(p,q)}. \end{aligned} \quad (62)$$

By Theorem 4.10, $\|g\|_{L(p,q)} \sim \|g\|_{L_{w(p,q)}^q}$. By Lemma 5.6, $Tg \in GM^+$, hence $\|Tg\|_{L(p,q)} \sim \|Tg\|_{L_{w(p,q)}^q}$. Therefore, (62) can be written as

$$\begin{aligned} \left(\int_0^\infty x^{\frac{q}{p}} \left(\frac{1}{x} \int_0^x g(t) dt \right)^q \frac{dx}{x} \right)^{\frac{1}{q}} &\sim \|Tg\|_{L(p,q)} \\ &\leq C(p, q) \|g\|_{L(p,q)} \sim C(p, q) \left(\int_0^\infty x^{\frac{q}{p}} (g(x))^q \frac{dx}{x} \right)^{\frac{1}{q}}. \end{aligned}$$

Since $g(t) = \frac{f(t)}{t}$, we obtain

$$\left(\int_0^\infty x^{\frac{q}{p}-q} \left(\int_0^x f(t) \frac{dt}{t} \right)^q \frac{dx}{x} \right)^{\frac{1}{q}} \leq C(p, q) \left(\int_0^\infty x^{\frac{q}{p}-q} (f(x))^q \frac{dx}{x} \right)^{\frac{1}{q}}. \quad (63)$$

Inequality (61) for $\alpha \leq 1$ follows from (63) if we take $\alpha = 1 - \frac{1}{p} = \frac{1}{p'}$.

2) Consider $\alpha > 1$. Take $g(t) = t^{\epsilon-\alpha} f(t)$, where $0 < \epsilon < 1$. Clearly, $g \in GM$. Also:

$$x^{-\alpha} \int_0^x f(t) \frac{dt}{t} = x^{-\epsilon} \cdot \int_0^x x^{\epsilon-\alpha} f(t) \frac{dt}{t} \leq x^{-\epsilon} \cdot \int_0^x g(t) \frac{dt}{t}.$$

By (61), for $\epsilon < 1$ and $g \in GM$,

$$\begin{aligned} \left(\int_0^\infty \left(x^{-\epsilon} \int_0^x g(t) \frac{dt}{t} \right)^q \frac{dx}{x} \right)^{\frac{1}{q}} &\leq C(\alpha, q) \left(\int_0^\infty (x^{-\epsilon} g(x))^q \frac{dx}{x} \right)^{\frac{1}{q}} \\ &= C(\alpha, q) \left(\int_0^\infty (x^{-\alpha} f(x))^q \frac{dx}{x} \right)^{\frac{1}{q}}, \end{aligned}$$

and (61) for $\alpha > 1$ follows. $\square$

References.

- [1] Askey, R.; Wainger, S. Integrability theorems for Fourier series. *Duke Math. J.* 33, 223–228, 1966.
- [2] Bergh, J.; Löfström, J. Interpolation spaces. An introduction. *Grundlehren der Mathematischen Wissenschaften*, No. 223. Springer-Verlag, Berlin-New York, 1976. x+207 pp.
- [3] Booton, B. General monotone sequences and trigonometric series. *Math. Nachr.* 1-12, 2013.
- [4] Brudnyi, Yu. A.; Krugljak, N. Ya. Interpolation functors and interpolation spaces. Vol. I. North-Holland Mathematical Library, 47. North-Holland Publishing Co., Amsterdam, 1991. xvi+718 pp.
- [5] Gilbert, J. E. Interpolation between weighted L^p -spaces. *Ark. Mat.* 10, 235–249, 1972.
- [6] Hardy, G.H.; Littlewood, J.E. Some new properties of Fourier constants. *J.L.M.S.* 6, 3-9, 1931.
- [7] Hardy, G.H.; Littlewood, J.E. Some new properties of Fourier constants. *M.A.* 97, 159-209, 1926.
- [8] Hardy, G.H.; Littlewood, J.E.; Pólya, G. *Inequalities*. 2d ed. Cambridge, at the University Press, 1952.
- [9] Lifyand, E.; Tikhonov, S. A concept of general monotonicity and applications. *Math. Nachr.* 284, no. 8-9, 1083–1098, 2011.
- [10] Paley, R.E.A.C. "Some theorems on orthogonal functions (1)." *Studia Mathematica* 3.1: 226-238, 1931.
- [11] Sagher, Y. An application of interpolation theory to Fourier series. *Studia Math.* 41, 169–181, 1972.
- [12] Sagher, Y. Integrability conditions for the Fourier transform. *J. Math. Anal. Appl.* 54, no. 1, 151–156, 1976.
- [13] Shah, S. M. A note on quasi-monotone series. *Math. Student* 15 (1947), 19–24, 1948.
- [14] Tikhonov, S. Trigonometric series with general monotone coefficients. *J. Math. Anal. Appl.* 326, no. 1, 721–735, 2007.
- [15] Zygmund, A. *Trigonometric series*. 2nd ed. Vols. I, II. Cambridge University Press, New York 1959. Vol. I. xii+383 pp.; Vol. II. vii+354 pp.